

# Fresubin® YODRINK

Available in three delicious flavours  
in 200 ml EasyBottle



## Nutritional Information

### Nutrition values

Flavour Raspberry		per 100 ml	bottle = 200 ml
<b>Energy</b>	kJ (kcal)	630 (150)	1260 (300)
<b>Fat</b>	g	4,7	9,4
of which saturates	g	0,6	1,2
of which mono-unsaturates	g	3,1	6,2
of which polyunsaturates	g	1,0	2,0
<b>Carbohydrate</b>	g	20	40
of which sugars	g	15	30
of which lactose	g	3,5	7,0
<b>Fibre</b>	g	0,2	0,4
<b>Protein</b>	g	7,5	15,0
<b>Salt</b>	g	0,15	0,30
<b>Vitamins</b>			
Vitamin A	µg RE <sup>o</sup>	190	380
Vitamin D	µg	3,8	7,6
Vitamin E	mg α-TE <sup>oo</sup>	4,0	8,0
Vitamin K	µg	20	40
Vitamin C	mg	19	38
Thiamin	mg	0,3	0,6
Riboflavin	mg	0,5	1,0
Niacin	mg/mg NE <sup>ooo</sup>	1,5/3,0	3,0/6,0
Vitamin B <sub>6</sub>	mg	0,4	0,8
Folic acid	µg	62,5	125
Vitamin B <sub>12</sub>	µg	0,8	1,6
Biotin	µg	9,4	18,8
Pantothenic acid	mg	1,1	2,2
<b>Minerals, trace elements and other<sup>#</sup> nutrients</b>			
Sodium	mg	60	120
Potassium	mg	165	330
Chloride	mg	85	170
Calcium	mg	240	480
Phosphorus	mg	95	190
Magnesium	mg	18	36
Iron	mg	2,5	5,0
Zinc	mg	2,0	4,0
Copper	mg	0,4	0,8
Manganese	mg	0,5	1,0
Fluoride	mg	0,3	0,6
Selenium	µg	14	28
Chromium	µg	13	26
Molybdenum	µg	19	38
Iodine	µg	37,5	75,0
Choline <sup>#</sup>	mg	32	64
<b>Osmolarity</b>	mosmol/l	885	
<b>Water</b>	ml	75	150

### Caloric distribution (kJ%)

Fat 28, carbohydrate 51,7, fibre 0,3, protein 20

<sup>o</sup>retinol equivalents (RE), <sup>oo</sup>alpha-tocopherol equivalents (α-TE), <sup>ooo</sup>niacin equivalents (NE)

## General Information

### Food for special medical purposes.

For the dietary management in case/risk of malnutrition, esp. with increased energy and protein needs. Nutritionally complete, high energy (1,5 kcal/ml), high protein oral nutritional supplement. High in calcium and vitamin D, contains lactose, gluten free.

### Dosage:

2-3 bottles/day for supplementary nutrition or 5 bottles/day for complete nutrition, or as advised by a healthcare professional.

### Important notice:

To be used under medical supervision. Suitable as sole source of nutrition. Not suitable for children < 3 years. Use with caution in children < 6 years.

### Storage:

At room temperature. Once opened refrigerate and use within 24 hours.

### Instructions for use:

Best served chilled. Shake well before use.  
Packaged in a protective atmosphere.

### Additional Considerations:

Not suitable whenever enteral nutrition is not permitted such as in acute gastrointestinal bleeding, ileus and shock. Use with caution in severe organ failure with impaired metabolism and severe forms of malassimilation. Not suitable for patients with congenital inability to metabolise nutrients contained in Fresubin YODRINK.

## Ingredients

**Ingredients (Flavour Raspberry):** Water, sucrose, whey protein, skimmed milk yoghurt powder, vegetable oils (rapeseed oil, sunflower oil), maltodextrin, acidity regulator (E 270), calcium lactate, emulsifiers (E 471, soya lecithins), thickener (E 440), colour (E 120), flavourings, choline bitartrate, vitamin C, sodium chloride, ferric diphosphate, magnesium oxide, zinc sulphate, vitamin E, pantothenic acid, manganese chloride, niacin, cupric sulphate, vitamin B<sub>6</sub>, sodium fluoride, riboflavin, vitamin A, thiamin, folic acid, chromium chloride, potassium iodide, sodium selenite, vitamin K, sodium molybdate, biotin, vitamin D, vitamin B<sub>12</sub>.

Nutrition information for flavour Raspberry. Detailed nutrition information of the other flavours can be found at [www.fresubin.com](http://www.fresubin.com)

## Flavours

Apricot-Peach, Lemon, Raspberry

- 1) Catwood AL, Elia M, Stratton RJ. Systematic review and meta-analysis of the effects of high protein oral nutritional supplements. *Ageing Res Rev*. 2012;11(2):278-296. 2) Dent E, Morley JE, Cruz-Jentoft AJ, Woodhouse L, Rodriguez-Manas L, Fried LP, et al. Physical Frailty: ICFSR International Clinical Practice Guidelines for Identification and Management. *J Nutr Health Aging*. 2019;23(9):771-87. 3) Dent E, Lien C, Lim WS, Wong WC, Ng TP, et al. The Asia-Pacific Clinical Practice Guidelines for the Management of Frailty. *J Am Med Dir Assoc*. 2017;18(7):564-75. 4) Morley JE, Vellas B, van Kan GA, Anker SD, Bauer JM, Bernabei R, et al. Frailty consensus: a call to action. *J Am Med Dir Assoc*. 2013;14(6):392-7. 5) Bischoff-Ferrari HA, Willett WC, Oran EJ, et al. Pooled analysis of vitamin D dose requirements for fracture prevention. *N Engl J Med*. 2012; 367(1): 40- 49. 6) Bischoff-Ferrari HA, Dawson-Hughes B, Staehelin HB, et al. Fall prevention with supplemental and active forms of vitamin D: a meta-analysis of randomised controlled trials. *The BMJ*. 2009; 339: b3692. 7) Dawson-Hughes B, Mithal A, Bonjour JP, et al. IOF position statement: vitamin D recommendations for older adults. *Osteoporos Int*. 2010; 21(7): 1151-1154. 8) Tang BM, Eslick GD, Nowson C, Smith C, Bousoussan A. Use of calcium or calcium in combination with vitamin D supplementation to prevent fractures and bone loss in people aged 50 years and older: a meta-analysis. *Lancet*. 2007; 370(9588): 657-664. 9) Larsen ER, Moskilde L, Foldspang A. Vitamin D and calcium supplementation prevents osteoporotic fractures in elderly community dwelling residents: a pragmatic population-based 3-year intervention study. *J Bone Miner Res*. 2004; 19(3): 370-378. 10) Hubbard GP, Elia M, Holdaway A, et al. A systematic review of compliance to oral nutritional supplements. *Clin Nutr*. 2012;31(3): 293-312.



**FRESENIUS  
KABI**  
caring for life

Fresenius Kabi Deutschland GmbH  
61346 Bad Homburg, Germany  
Phone: +49 (0) 61 72 / 686-0  
[Enteral.nutrition@fresenius-kabi.com](mailto:Enteral.nutrition@fresenius-kabi.com)  
[www.fresenius-kabi.com](http://www.fresenius-kabi.com)